

Clinical Efficacy of Low Carbohydrate Diet (LCD) For Detail Analysis of Daily Meal and Blood Glucose

Urasaki H^{1,2}, Bando H^{2,3,4,*}, Urasaki H², Hayashi Y², Ogawa H^{2,4} and Nagahiro S⁴

¹Kainan Hospital, Naruto city, Tokushima, Japan

²Tokushima Low Carbohydrate Diet Promotion Association, Tokushima, Japan.

³Medical Research, Tokushima University, Tokushima, Japan

⁴Yoshinogawa Hospital, Tokushima, Japan

⁵Emeritus Professor, Former Director of Tokushima University Hospital, Tokushima, Japan

*Corresponding author: Bando H, Tokushima University, Medical Research, Nakashowa 1-61, Tokushima 770-0943 Japan; Tel: +81-90-3187-2485; E-mail: pianomed@bronze.ocn.ne.jp

Received date: 19 December 2021; Accepted date: 22 December 2021; Published date: 29 December 2021

Citation: Urasaki H, Bando H, Urasaki H, Hayashi Y, Ogawa H, Nagahiro S (2021). Clinical Efficacy of Low Carbohydrate Diet (LCD) For Detail Analysis of Daily Meal and Blood Glucose. SunText Rev Med Clin Res 2(4): 144.

DOI: <https://doi.org/10.51737/2766-4813.2021.044>

Copyright: © 2021 Urasaki H, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background: Background: Regarding nutritional therapy for obesity and diabetes, low carbohydrate diet (LCD) has been effective.

Case presentation: The case is 61-year-old male with obesity and type 2 diabetes mellitus (T2DM). He has been an athlete for long, and exercised much in hot climate during Mar-Sept, 2021. He often drank much amount of oral rehydration solution (ORS) and sports drink solution (SDS). In late Sept, he felt strong fatigue, weight loss, thirsty, and visited a clinic. He was diagnosed to have severe T2DM with HbA1c 14.0%.

Results: He started to take super-LCD and to take pictures of all meals every day. He can continue them perfectly and HbA1c decreased to 10.7% in October and 7.8% in November. He has checked pre-prandial and post-prandial blood glucose, ranging almost 120-200 mg/dL due to LCD continuation. Daily carbohydrate intake amounts were checked, which were 87g, 137g, 81g and 136g.

Discussion and Conclusion: His intake of ORS and SDS seemed to be adequate judging from their difference. The clinical course was successful, that was probably due to satisfactory lifestyle, continuing LCD and recording diet method, accumulating picture data of all meal, calculating carbohydrate and calorie intake every day. This case become a significant reference.

Keywords: Low carbohydrate diet (LCD); Type 2 diabetes mellitus (T2DM); Oral rehydration solution (ORS); Sports drink solution (SDS); Daily carbohydrate intake; Recording diet

Introduction

Diabetes mellitus (DM) has been gradually increasing for long years worldwide, reaching to the degree of pandemic. It includes apparent DM and also undiagnosed diabetes mellitus (UDM). International Diabetes Federation (IDF) has continued various management for DM and presented the estimates of UDM [1]. From worldwide data, approximately one in two adult population (20 to 79 years) with diabetes were unaware of their diabetes status, which was 239.7 million and 44.7%. Combined of clinical DM and UDM, the prevalence was reported to be 9.3% in 2019, and will be up to 10.2% and 10.9% in 2030 and 2045,

respectively [2]. Obesity is usually the basis of diabetes and metabolic syndrome [3]. Therefore, it is necessary to take appropriate measures against obesity from an early stage. Proper diet and exercise are fundamentally essential for these responses [4]. For nutritional therapy for obesity and diabetes, low carbohydrate diet (LCD) has been effective. Before LCD, calorie restriction (CR) was standard diet method by the calculation of total calorie a day. However, several researchers in western countries reported clinical efficacy of LCD from historical point of view [5]. After that, detail comparison of three meals with LCD, CR and Mediterranean diet was reported [6]. Recently, LCD has been estimated to become first approach for the

management of diabetes and obesity [7]. Similar to European countries, LCD in Japan was initiated by authors and collaborators [8]. For practical application of LCD for everyone, three types of LCD were presented, which are super LCD, standard LCD and petite LCD. They are easy to understand to continue for daily diet therapy, where carbohydrate content by calorie ratio is 12%, 26% and 40%, respectively [9]. Successively, we have continued the development of LCD in the light of medical and social aspects by establishment of Japan Low Carbohydrate Diet Promotion Association (JLCDPA) [10]. We have treated thousands of obesity patients for LCD, in which 25% of the patients achieved reduction more than 10% of former body weight [11]. Authors and collaborators continued practice and research of diabetes and LCD for years [12]. Furthermore, clinical research of insulin secretion for meal tolerance test (MTT) was conducted [13]. During our medical practice, 61-year-old male with obesity and type 2 diabetes mellitus (T2DM) was treated for LCD. The case showed remarkable improvement of diabetic status by strict LCD meal with complete recording data. His clinical course with some perspectives will be described in this article.

Case Presentation

History of present illness (HPI)

The case is a 61-year-old man with obesity and type 2 diabetes mellitus (T2DM). When he was a child, he started to play baseball. He became an excellent pitcher and he could throw fast ball about 150 km/hr. During senior high school, he was known widely and had the possibility to be selected to the draft selection of Nippon Professional Baseball (NPB) Organization. However, he had to throw too much games, and then he hurt his shoulder and elbow. He had to resign entering NPB. After that, he has worked in the administration section of the hospital. With his diligence and trust from others, he became the general manager of the hospital and developed his career in the medicinal practice. Furthermore, he contributed much in the baseball and softball sports fields. He won in several baseball tournament, including all-Japan national sports festival as a player, coach and manager. He also continues teaching other coaches in senior baseball teams and developing research of sports medicine in Research association of baseball Tokushima (RABT).

Concerning his health status, he was healthy until 30's. During his 40's, his weight increased from 78kg to more than 100kg. He was diagnosed as obesity and T2DM in his late 50's. Then, he was advised to continue LCD, resulting in 10kg weight reduction until 2020. When he played many softball games in Mar-Sept, 2021, he drank much amount of oral rehydration solution (ORS) and sports drink solution (SDS). He felt strong fatigue and weight loss in late Sept, and his HbA1c was proved to be 14.0%.

Physical and Other Examination

As to physical status, his consciousness, speech and behaviour were normal. Vitals were pulse rate 56/min, BP 136/82 with normal temperature and respiration. His physique was 181 cm, 87kg, 26.6 kg/m² in BMI and 88 cm in abdominal circumference. For reference, his former physique during 50's was 104kg, 31.7 kg/m² and 110cm when he was 50-56 years old. His physical examination showed unremarkable in the lung, heart, abdomen and neurological findings. The results of several examinations in Sept, 2021 were as follows: Hb 16.2 g/dL, RBC 508 x 10⁶ /μL, WBC8600 /μL, Plt 21.1 x 10⁴ /μL, BUN 14 mg/dL, Cr 0.7 mg/dL, uric acid 3.0 mg/dL, TG 100 mg/dL, HDL-C 65 mg/dL, LDL-C 142 mg/dL, TP 7.3 g/dL, Alb 4.3 g/dL, AST 22 U/L, ALT 28 U/L, GGT 25 U/L, ALP 193 U/L (100-340), LDH 141 U/L (120-240), CRP 0.02 mg/dL, Concerning diabetes, HbA1c 14.0%, post-prandial blood glucose 292 mg/dL. Other examinations were as follows: Chest X-P was unremarkable, and electrocardiogram (ECG) revealed within normal limits. The examination of pulse wave velocity (PWV) showed that ankle brachial index (ABI) 1.19/1.19, cardio-ankle vascular index (CAVI) 9.3/9.2 in right/left, respectively.

Clinical Course and Results

His severe diabetic status was found in later September, 2021. At once, he was to begin LCD meals for three meals a day. He could continue LCD every day and the actual meals are recorded by camera. Furthermore, he could check the values of blood glucose frequently, which were regularly connected to pre-prandial and post-prandial timing. The results of fasting, and 45 min postprandial data of lunch and supper are summarized (Table 1).

The values of pre-prandial and post-prandial glucose were well controlled between about 120-200 mg/dL, which was due to limited intake of carbohydrate. Furthermore, his daily meals are recorded every day. The actual meals on Oct 20, Nov 5, Nov 12 and Nov 19 are observed (Figure 1).

Taken food included meat, salad, soup, Japanese style pot, bread, noodle, and so on. They showed less carbohydrate intake amount per day, which were 87g, 137g, 81g and 136g, respectively. For one of 4 days in Figure 1, the detail content of the meal on Oct 2 are as follows: i) breakfast: low-carbohydrate chocolate bread 1.7g, cream stew-style soup 12g, coffee 1g, ii) lunch: lunch box including mackerel dishes 6.5g, roast beef 7.7g, salad 1g, miso soup 13g, iii) supper: omelette 27.7g, salad 1g, boiled seaweed 7.6g, iv) snack: fruit citrus 7.4g. For total calculation, carbohydrate amount is 87g/day, and total calories is 1589 kcal/day. By continuing LCD, his HbA1c levels were acutely decreased. During Sept-Nov, 2021, HbA1c values were 14.0%, 10.3% and 7.8%, respectively.



Figure 1: Actual meal for LCD with carbohydrate amount.

Discussion

In current case, some characteristic points are observed in the following: i) he has been an athlete working in the hospital, who has rather enough knowledge for general medicine and sports medicine, ii) he was treated obesity and T2DM by LCD about a year ago, iii) drinking much amount of oral rehydration solution (ORS) and sports drink solution (SDS) can become a trigger for exacerbation of diabetic control, iv) continuation of LCD and recording every meal was effective for clinical improvement. Their perspectives are described for the discussion in order. Firstly, the case has been a baseball player so far, and was not obese until 40 years old. During 40s and 50s, he came to have obesity and T2DM. He tried some kinds of nutritional and exercise approaches for weight reduction, including calorie restriction (CR) and resistance muscle training. However, they were not effective or successful [14].

Table 1: The results of fasting, and 45 min postprandial data of lunch and supper are summarized.

Month	Date	breakfast-0min	breakfast-45min	supper-45min	carbohydrate	calorie	actual
		blood glucose	blood glucose	blood glucose	amount	energy	meal
		(mg/dL)	(mg/dL)	(mg/dL)	(g/day)	(kcal/day)	(Fig 1)
Oct	4		121		66	1882	
	6	148			50	1787	
	8			130	140	1573	
	11		205		101	1313	
	13	111			104	2387	
	15			220	91	1490	
	18		226		83	1348	
	20	118			87	1589	Fig 1a
	22			133	99	1970	
Nov	1		132		176	1400	
	3	132			130	1633	
	5			166	137	1663	Fig.1b
	8		188		110	1970	
	10	121			150	1679	
	12			150	81	1494	Fig 1c
	15		154		132	2032	
	17	123			103	2294	
	19			145	136	1182	Fig 1d
	22		143		166	1839	

From current episode, he came to know actual efficacy of LCD and recording diet method [15]. Secondly, he had the experience for LCD a year ago. At that time, clinical efficacy was not enough. The reason would be the degree of carbohydrate restriction. We have developed LCD movement medically and socially through Japan LCD Promotion Association (JLCDPA) [9]. There are three levels of LCD, which are super LCD, standard LCD and petite LCD, associated with carbohydrate ratio

of 12%, 26% and 40%, respectively. He applied petite LCD in autumn 2020, and super-LCD in autumn 2021 [16]. Authors and colleagues have continued LCD treatment for more than 2700 cases with diabetes, obesity and metabolic syndrome [17]. The results showed that weight reduction was 6.6% in average, and 2.6%-9.8% in the quartiles of 25%-75% associated with mean weight reduction as 4.3 kg. Thirdly, he drank much amount of ORS and SDS during Mar-Sept 2021, and this episode may bring

the exacerbation of T2DM. Actually, he drank about 1000cc of ORS in Mar-May with rather mild climate period. After that he drank about 2000-3000cc of SDS in Jun-Sept with severe hot climate period. Thus, his method of ingesting water seemed to be appropriate. The reason is the difference in the components of ORS and SDS [18]. The purpose of ORS is to treat dehydration caused by illness and diarrhea. On the other hand, the purpose of SDS is to replenish water, sugar, and minerals due to a large amount of sweating in sports [19]. However, these objectives have been often confused in our usual daily lives [20].

As to ORS, he utilized Pocari sweat (Otsuka Pharma, Japan) [21]. The content shows carbohydrate 6.2g, protein and fat 0g, energy 25kcal, NaCl 0.12g, K 20mg, Ca 2mg, Mg 0.6mg. For SDS, he used Body mainte (Otsuka Pharma, Japan) [22]. The content was carbohydrate 4.4g, protein and fat 0g, Energy 18kcal, NaCl 0.13g, K 20mg, Ca 2mg, Mg 0.6mg, as well as lactic acid bacterium B240. B240 was found from a plant, which has been used for fermented tea in Thailand [23]. From immunological point of view, B240 (*Lactobacillus pentosus* ONRICb0240) has been utilized for developing new conditioning beverage [22]. Several beneficial enhancements have been observed including blood T-helper, NK cell, IgA/IgG levels, and other immune functions (Kiso). Some positive efficacy would be found for elderly people, physical exercise [24-26]. Fourthly, continuation of LCD with recording data has successfully brought him clinical improvement for glucose variability. Formerly, American Diabetes Association (ADA) presented the official comment, in which only carbohydrate can directly affect blood glucose levels [27]. It is well-known that diabetic pathophysiological problems are due to elevated blood glucose, such as microvascular and macrovascular disorders [28]. Then, restriction of carbohydrate intake per os can refrain from hyperglycemia. Several evidence of LCD efficacy were reported from various studies [6,29]. Further beneficial data for LCD were found in some mega-studies [30,31]. From these reports, carbohydrate loading per os may become one of the evaluation methods. Meal Tolerance Test (MTT) is a useful way to apply certain amount of carbohydrate for breakfast. Authors have proposed MTT using 70g-Carbo breakfast [32]. MTT seems to be applied more for simple examination for glucose variability [33]. Regarding most recent reports on LCD, significant tendency can be found for short and long period [34], and for LCD and very LCD. Future comparative study for LCD seems to focus in detail differences, with expectation for beneficial diet method for diabetes and obesity [35]. In summary, this case report showed a diabetic male with obesity and T2DM. Several problems and discussion were described. There are some limitations such as detail situation of diabetes, hydration, ORS, SDS, and actual LCD meals, which will be further investigated and follow up. We hope that this meaningful case will be useful for medical development of diabetes and research in the future.

Ethical Considerations

Current study was performed in compliance with the adequate ethical principles. It is due to the Declaration of Helsinki. Moreover, some commentary was shown for the Ethical Guidelines for Research in the medical field for Human beings and in the conduction of the Good Clinical Practice (GCP). Authors have been provided written informed consent from the patient himself in this study. In addition, authors established the ethical committee for the clinical research including medical, pharmacological and legal specialties. We have fully discussed the detail of the study and made confirmation that this study would be proper and agreed with all members.

Conflict of Interest

The authors declare no conflict of interest.

Funding

There was no funding received for this paper.

References

1. Ogurtsova K, Guariguata L, Barengo NC, Ruiz PLD, Sacre JW, Karuranga S, et al. IDF Diabetes Atlas: Global estimates of undiagnosed diabetes in adults for 2021. *Diabetes Res Clin Pract.* 2021; 1: 109118.
2. Saeedi P, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, et al. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas. *Diab Res Clin Pract.* 2019; 157: 1-10.
3. Chobot A, Kowolik KG, Sokołowska M, Chobot PJ. Obesity and diabetes-Not only a simple link between two epidemics. *Diabetes Metab Res Rev.* 2018; 34.
4. Facilitating behavior change and wellbeing to improve health outcomes: Standards of medical care in diabetes. *Diabetes Care.* 2021; 44: 53-72.
5. Macedo RCO, Santos HO, Tinsley GM, Oliveira AR. Low-carbohydrate diets: Effects on metabolism and exercise - A comprehensive literature review. *Clin Nutr ESPEN.* 2020; 40: 17-26.
6. Shai I, Schwarzfuchs D, Henkin Y. Dietary Intervention Randomized Controlled Trial (DIRECT) Group. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med.* 2008; 359: 229-241.
7. Unwin D, Unwin J, Crocombe D, Delon C, Guess N, Wong C, et al. Renal function in patients following a low carbohydrate diet for type 2 diabetes: a review of the literature and analysis of routine clinical data from a primary care service over 7 years. *Curr Opin Endocrinol Diabetes Obes.* 2021; 28: 469-479.
8. Ebe K, Ebe Y, Yokota S, Matsumoto T, Hashimoto M, Sakai Y, et al. Low Carbohydrate diet (LCD) treated for three cases as diabetic diet therapy. *Kyoto Med Association J.* 2004; 51: 125-129.

9. Bando H. Useful tips for actual low carbohydrate diet (LCD) with super-, standard- and petite-LCD methods. *EC Nutrition*. 2020; 15: 1-4.
10. Muneta T, Kagaguchi E, Nagai Y, Matsumoto M, Ebe K, Watanabe H, et al. Ketone body elevation in placenta, umbilical cord, newborn and mother in normal delivery. *Glycat Stress Res*. 2016; 3: 133-140.
11. Nakamura T, Kawashima T, Dobashi M, Narita A, Bando H. Effective nutritional guidance for obesity by Low Carbohydrate Diet (LCD). *Asp Biomed Clin Case Rep*. 2019; 2: 16-21.
12. Kanazawa S, Kato Y, Bando H. Significant weight reduction by super low carbohydrate diet with improved lipid profiles. *Int Med*. 2020; 2: 175-178.
13. Ebe K, Hashimoto M, Bando H, Bando M, Muneta T. Proposal of Meal Tolerance Test (MTT) For investigating ability of insulin secretion for small carbohydrate load. *Diab Res Open Access*. 2020; 2: 31-37.
14. Li S, Ding L, Xiao X. Comparing the efficacy and safety of low-carbohydrate diets with low-fat diets for type 2 diabetes mellitus patients: A systematic review and meta-analysis of randomized clinical trials. *Int J Endocrinol*. 2021; 2021: 8521756.
15. Nakamura T, Bando H, Kawashima T, Dobashi M, Narita A. Weight reduction by effective protocol of diet and exercise. *J Diab Obes Metab*. 2019; 2: 106.
16. Bando H, Ebe K, Muneta T, Bando M, Yonei Y. Clinical effect of low carbohydrate diet (LCD): Case Report. *Diabetes Case Rep*. 2017; 2: 124.
17. Schleh MW, Charles L, Dumke CL. Comparison of sports drink versus oral rehydration solution during exercise in the heat. *Wildern Environment Med*. 2018; 29: 185-193.
18. Hannon MP, Close GL, Morton JP. Energy and macronutrient considerations for young athletes. *Strength Condit J*. 2020; 42: 109-119.
19. Orrù S, Imperlini E, Nigro E, Alfieri A, Cevenini A, Polito R, et al. Role of functional beverages on sport performance and recovery. *Nutrients*. 2018; 10: 1470.
20. Dana TGP, Pramudana KAS. The role of brand image in mediating the influence of product quality on re-purchase intentions pocari sweat isotonic beverages. *Am J Human Soc Sci Res*. 2021; 5: 255-258.
21. Body Mainte: <https://www.otsuka.co.jp/bdm/products/>
22. Lee M, Kim K. Innovative and practical conditioning beverages for public health and athletic performance: Focus on immunopotentiality by lactic acid bacteria B240. *J Exerc Nutrition Biochem*. 2019; 23: 13-15.
23. Kiso M, Takano R, Sakabe S, Katsura H, Shinya K, Uraki R, et al. Protective efficacy of orally administered, heat-killed *Lactobacillus pentosus* b240 against influenza A virus. *Sci Rep*. 2013; 3: 1563.
24. Shinkai S, Toba M, Saito T, Sato I, Tsubouchi M, Taira K, et al. Immunoprotective effects of oral intake of heat-killed *Lactobacillus pentosus* strain b240 in elderly adults: a randomised, double-blind, placebo-controlled trial - CORRIGENDUM. *Br J Nutr*. 2016; 116: 1138-1140.
25. Shimizu K, Sato H, Suga Y, Yamahira S, Toba M, Hamuro K, et al. The effects of *Lactobacillus pentosus* strain b240 and appropriate physical training on salivary secretory IgA levels in elderly adults with low physical fitness: a randomized, double-blind, placebo-controlled trial. *J Clin Biochem Nutr*. 2014; 54: 61-66.
26. American Diabetes Society. Nutrition principles and recommendations in diabetes. *Diabetes Care*. 2004; 27: 36.
27. Mauricio D, Alonso N, Gratacos M. Chronic diabetes complications: The need to move beyond classical concepts. *Trends Endocrinol Metab*. 2020; 31: 287-295.
28. Shai I, Schwarzfuchs D, Henkin Y. Dietary intervention randomized controlled trial (DIRECT) group. Weight loss with a low-carbohydrate, Mediterranean, or Low-Fat Diet. *N Engl J Med*. 2008; 359: 229-241.
29. Feinman RD, Pogozelski WK, Astrup A, Bernstein RK, Fine EJ, Westman EC, et al. Dietary carbohydrate restriction as the first approach in diabetes management: Critical review and evidence base. *Nutrition*. 2015; 31: 1-13.
30. Meng Y, Bai H, Wang S, Li Z, Wang Q, Chen L, et al. Efficacy of low carbohydrate diet for type 2 diabetes mellitus management: A systematic review and meta-analysis of randomized controlled trials. *Diabetes Res Clin Pract*. 2017; 131: 124-131.
31. Oh R, Uppaluri KR. Low carbohydrate diet. In: StatPearls. Treasure Island (FL): StatPearls Publishing. 2020.
32. Bando H, Ebe K, Muneta T, Bando M, Yonei Y. Proposal for Insulinogenic Index (IGI)-Carbo70 as experimental evaluation for diabetes. *J Clin Exp Endocrinol*. 2017; 1: 102.
33. Ebe K, Hashimoto M, Bando H, Bando M, Muneta T. Proposal of Meal Tolerance Test (MTT) for investigating ability of insulin secretion for small carbohydrate load. *Diab Res Open Access*. 2021; 2: 31-37.
34. Barber TM, Hanson P, Kabisch S, Pfeiffer AFH, Weickert MO. The Low-Carbohydrate Diet: short-term metabolic efficacy versus longer-term limitations. *Nutrients*. 2021; 13: 1187.
35. Goldenberg JZ, Day A, Brinkworth GD, Sato J, Yamada S, Jonsson T, et al. Efficacy and safety of low and very low carbohydrate diets for type 2 diabetes remission: systematic review and meta-analysis of published and unpublished randomized trial data. *BMJ*. 2021; 372: 4743.