



# Helicobacter Pylori Infection Association with Anxiety in Patients with Upper Gastrointestinal Symptoms in Hadhramout

Daakeek AM<sup>1\*</sup>, Musiaan NS<sup>2</sup> and Al-Zaazaai AA<sup>2</sup>

<sup>1</sup>Associated Professor of Medicine, Medical Department, Faculty of Medicine and Health Sciences, Hadhramout University, Mukalla, Yemen

<sup>2</sup>Department of Pharmacy, Wenzhou Medical University, Wenzhou, PR, China

\*Corresponding author: Dr. Ahmed Mubarak Daakeek, MD, Associated Professor of Medicine, Medical Department, Faculty of Medicine and Health Sciences, Hadhramout University, Mukalla, Yemen; E-mail: daakeek@gmail.com [daakeek@hotmail.com](mailto:daakeek@hotmail.com)

Received date: 31 October 2023; Accepted date: 15 November 2023; Published date: 26 November 2023

Citation: Daakeek AM, Musiaan NS, Al-Zaazaai AA (2023) Helicobacter Pylori Infection Association with Anxiety in Patients with Upper Gastrointestinal Symptoms in Hadhramout. SunText Rev Med Clin Res 4(4): 191.

DOI: <https://doi.org/10.51737/2766-4813.2023.091>

Copyright: © 2023 Daakeek AM, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

## Abstract

**Background:** Helicobacter pylori (H. pylori) infection is highly prevalent infection in Yemen that has been linked with numerous gastrointestinal diseases. An association between it and anxiety has been observed. Few studies have focused on the relationship between H. pylori infection with anxiety.

**Objective:** In this study, we aimed to assess the relation between anxiety and H. pylori infection in patients with gastrointestinal symptoms.

**Material and methods:** This prospective study included 233 adult patients with upper gastrointestinal symptoms who were presented to medical outpatient in Mukalla city between August 2022 and February 2023. Data collected using four sections perform: demographic features, upper gastrointestinal manifestations, anxiety assessment using the Hamilton Anxiety Scale (HAS), and H.pylori detection. Patients were divided into three groups based on their age and two groups based on H.pylori status.

**Results:** Most of the patients were female, with a female to male ratio of 2.6:1, and the mean age was 35.03 years (SD ± 13.732), with the 15-35 years age group being the most common (54%). Epigastric pain is a common upper gastrointestinal symptom, while fear and insomnia are also common anxiety symptoms. Most cases were positive for H;pylori ( 65.7%). Anxiety was observed in (96.1%), of H. pylori positive patients and most are moderately anxious (55.2%), with significant correlations between them (p-value 0.013).

**Conclusion:** H. pylori infection was common in female aged less than 55 years, and is significantly associated with anxiety.

**Keywords:** H. pylori infection; HAS score; Anxiety

## Introduction

Helicobacter (H. Pylori) is gram-negative and spiral, and has multiple flagella at one end, which make it motile, allowing it to burrow and live beneath the mucus layer adherent to the epithelial surface [1]. The prevalence of H. Pylori is high in developing countries (80–90% of the population) and much lower (20–50%) in developed countries. Infection rates are highest in lower-income groups. Infection is usually acquired in childhood; although the exact route is uncertain, it may be faecal–oral or oral–oral [2]. The vast majority of colonized people remain healthy and asymptomatic, and only a minority develop clinical

disease [1]. H. Pylori infection has been recognized as the main risk factor for many upper digestive tract disorders and complications, including life-threatening bleeding from the digestive tract, gastritis, non-ulcer dyspepsia, mucosa-associated lymphoid tissue (malt) lymphoma and gastric cancer [3]. The presence of h. Pylori-negative upper digestive tract abnormalities, such as peptic ulcers and the fact that not all h. Pylori-infected patients develop ulcers, with only 10%-15% of them presenting dyspeptic symptoms [3], suggest a role for other individual factors, including nervous system imbalance, as an indispensable cofactor in gastritis or ulcer disease pathogenesis [4]. These imply that h. Pylori infection may induce changes in the function and

morphology of the digestive tract both directly through cytotoxin release and inflammatory process activation, and indirectly, via the brain-gut axis [5]. Gut-brain axis is defined as the relationship of gastrointestinal with brain function and mental status. Mental stress is associated with personality-dependent gastric acid secretion changes [6]. Moreover, gastric inflammation leads to anxiety and depression-like behaviors via the neuroendocrine pathways especially in female [7]. There are several studies have reported that the change in the gut microorganisms could cause emotional disturbances, behavioral changes as well as mental health problems, but the mechanism is still under investigation [8]. This study aims to estimate the relations of H. pylori infections to the psychiatric manifestation in patients with upper gastrointestinal symptoms and the frequency of anxiety in those patients.

### Material and Methods

This study conducted in private medical center in mukalla city the capital of hadhramout governorate and the center of all eastern governors of yemen, which include shabowa, mahra and scotora besides hadhramout, and all this four governorates called hadhramout sector, so it is a reference city for many cases that come and transferred from these governors for treatment. This prospective study included 233 adult patients with upper gastrointestinal tract (ugit) symptoms who visited the medical outpatient between august 2022 and February 2023 (n = 223), examined the connection between H. Pylori infection and anxiety symptoms. a specially designed performa with four sections was used for data collection. Section one contains demographic details including age, gender and residence. All patients aged 15 years or older were included. The age is further grouped in decades to three age groups: 15–35, 36–55, and above 55 years.

The second section inquired about the details of a clinical profile including ugit symptoms which were defined at least two or more of the following symptom: epigastric pain or epigastric burning symptoms, retrosternal pain, heartburn, regurgitation, nausea and vomiting, early satiation, and postprandial fullness, and for the last 2 weeks. The third section consists of the hamilton anxiety scale (ha) assessment of anxiety, which is used to assess the severity of anxiety symptoms. The scale consists of 14 items; each item contains a number of symptoms, and each item is graded from zero to four according to severity. The results of the evaluation are reading as; mild severity =17 or less, moderate=18-24, and severe = 25-30. Fourth section records investigation primarily H.pylori infection by detection of stool antigen using rapid pylori antigen (trp) immunochromatography kit, as per manufacturer's instructions. The overall sensitivity is 97.6% and specificity is 96% [2]. The test is useful in the diagnosis of H. Pylori infection and for monitoring efficacy of eradication therapy; it is a cheap and easy method for the initial diagnosis of

H. Pylori infection. The patients were divided into two groups based on the presence or absence of h. Pylori infection: h. Pylori positive and h. Pylori negative groups. All cases were undergoing ultrasonographic imaging procedures to exclude biliary, pancreatic, and other diseases. Upper endoscopy was not done in almost all cases.




Figure 1: Frequency of age groups in all cases.




Figure 2: Distribution of cases according to age groups and sex.




Figure 3: Comparison of anxiety severity in Pylori positive and negative patients.

Data were analyzed using statistical software spss version 23, and microsoft office excel. Descriptive statistics of demographic variables were calculated including frequencies, percentages, means and ranges. Patient’s characters were compared using the chi square test. A1-sided p < 0.05 was considered statistically significant at the 95% confidence level.

### Ethical Consent

The college of medicine - Hadhrmout University (hucom), authorized the protocol for this study. All study participants gave their written consent after being informed of the objectives of our investigation.

### Results

The study included 233 patients with upper gastrointestinal symptoms, ranging in age from 15 to 73 years and mean age was 35.03 years (SD ±13.732), The majority of these patients were in age group of 15-35 years old (n=126, 54.1%), followed by age

group 36-55 years (n=82, 35.2%), and both constitute 89.3% of all cases, while older than 55 years age group considered the last group, as shown in Figure 1, and there is significant correlation between age and H.pylori status (p value 0.012). The majority of

cases being females (n=169, 72.5%) with female to male (ratio 2.6:1), the distribution of cases according to sex and age groups depicted in Figure 2.

**Table 1:** Anxiety symptom frequency in relation to H.pylori status.

Anxiety Items	H.pylori +ve		H.pylori -ve	
	No	%	No	%
Worries	115	80.4	52	77.6
Tension	88	61.5	32	47.8
Fears	132	93.3	61	91
Insomnia	129	90.2	60	89.6
Difficulty in concentration	88	61.5	33	49.3
Depressed mood	47	32.9	24	35.8
Somatic (muscular)	26	18.2	15	22.4
Somatic (sensory)	36	25.2	16	23.9
Cardiovascular symptoms	127	88.9	59	88.1
Respiratory symptoms	124	86.7	56	83.6
Gastrointestinal symptoms	123	86	54	80.6
Genitourinary symptoms	28	19.6	12	17.9
Autonomic symptoms	37	25.9	16	23.9
Behavior at interview	12	8.4	5	7.5

**Table 2:** Distribution of H. pylori cases according to age groups and sex.

Age group	H.pylori +ve			H.pylori -ve			Total cases
	Male	Female	Total	Male	Female	Total	
15-35 years	26	58	84	13	29	42	126
36-55 years	14	45	59	5	18	23	82
> 55 years	0	10	10	6	9	15	25
<b>Total</b>	40	113	153	24	56	80	233

Most of the cases from Hadhramout governorate (n=154, 66.1%), followed by Mahra governorate (n=47, 20.2%), Shabowa governorate (n=26, 11.2%) and Socotra governorate the last (n=6, 2.6%). All cases exhibit at least two symptoms of UGIT, and epigastric pain or epigastric burning symptoms are the most prevalent symptoms constitute 97.4% (n=227) of cases, while other symptoms are less frequent. Table 1 list the most frequent symptoms of anxiety according to HAS in relation to H.pylori status including fear, insomnia, cardiovascular, respiratory, and gastrointestinal symptoms which were the most common A majority of the cases (n=163, 65.7%) were H. pylori positive, with females accounting for the bulk of cases (n=113, 56%). Nearly half of the cases (51.3%) were in the 15–35 age range. Although there were 34.3% (n=80) H. pylori negative patients, the majority of them were female and belonged to the same age group. Table 2 displays the distribution of cases by age groups sex, and H. pylori status we employed the HAS questionnaire in the study to examine the level of anxiety in our cases, and we discovered that 214 of them (91.8%) had anxiety of varying

severity. Of the 147 individuals with H. pylori positivity who had anxiety status, n= 79, or 55.2%, had moderate anxiety (n=41, 27.9%) had mild anxiety, and (n=27, 18.4%) had severe anxiety. With a p value of 0.013, there is a significant association between anxiety and H. Pylori infection. In contrast, among the 80 cases in which H. Pylori negative , anxiety was present in 66 (82.5%) patients, the majority of whom had moderate intensity ((n=35, 43.8%), whereas moderate and severe anxiety were less common (22.5% and 17.5%, respectively). The level of anxiety in both H. pylori groups is shown in Figure 2.

### Discussion

H.pylori prevalence is higher in our country and is detected in patients with esophagitis, gastritis and peptic ulcer diseases as well as asymptomatic patients [9-11], and there is a relation between it and psychiatric manifestations in some studies [12,13]. In this study, we try to investigate the association between H.pylori and anxiety in 233 patients presents with UGIT manifestations. In this study, Hadhramout governorates accounted

for 66.1% of the cases. While Mahra, Shabow, and Scotra governments are less frequent, this is primarily because of their far distance from Mukalla, particularly Socotra, which is an island and has no other means of transport outside flights and boats, as well as their higher cost given the country current economic situation. The majority of *H. pylori* cases were females, which is consistent with research conducted in our country [9,10], as well as in Bahrain, Saudi Arabia, Egypt, Iran, and other nations in the Eastern Mediterranean region [5-16]. The age range 15-35 years was the most prominent and the most affected age group was under 55 years old, according to several research conducted in Yemen and its neighbouring countries [5-18]. No matter whether a patient had *H. pylori* or not, the most prevalent presenting symptom in almost all cases was epigastric pain resembling other studies [14]. While the most common symptoms of anxiety were fear and insomnia in 93.3% and 90.2% of all anxiety-related *H. pylori* positive patients, respectively. Our study demonstrates that patients with confirmed *H. pylori* infections have higher rates of anxiety than negative cases, with female patients being more likely to experience this. This finding is consistent with studies from Yemen [18], Bahrain [5], Saudi Arabia [16], and other countries that have also confirmed a connection between *H. pylori* and stress and anxiety [19-22]. The relationship between psychiatric disorders and *H. pylori* infection has been studied, and most patients in this research had some sort of anxiety with (26.8%) having mild Anxiety, (51.6%) moderate Anxiety, and (17.6%) severe Anxiety, while those without anxiety constitute about (3.9%) of all cases. This study finding was supported by some studies who discovered a strong positive relationship between *H. pylori* presence and anxiety symptoms [6, 20]. Additionally, numerous studies have shown that the elimination of *H. pylori* alleviates UGIT symptoms [23-25]. However, other research, like as the Kivrak, Y., et al. study, [26], disputes the relationship between the two.

*H. pylori* negative cases make up less percentage 34.3% than positive cases, there is considerable cases having anxiety, and this may attributable to infection and stress rather than *H. pylori* infection. *H. pylori* negative patients make up a smaller fraction (34.3%) than positive cases. There are many cases of anxiety for them, which may be caused by stress and fear of infection rather than *H. pylori* infection. In an intriguing approach, our research found that the positive *H. pylori* group had a highly significant rise in moderate and severe anxiety when compared to the negative *H. pylori* group.

## Conclusion

Hadhrumout and Yemen in general have a high rate of *H. pylori* infection, mostly in female patients under 55 years old, and a high rate of *H. pylori* infections associated with moderate to severe anxiety. Therefore, an evaluation of presence anxiety is crucial in

patients with gastrointestinal symptoms and symptoms of dyspepsia.

## References

1. Ralston SH, Penman ID, Strachan MW, Hobson R. Davidson's Principles and Practice of Medicine E-Book. Elsevier Health Sciences. 2018.
2. Kumar P, Clark M, Kumar and Clark's Clinical Medicine. 9<sup>th</sup> ed. Philadelphia, Pa.: Elsevier. 2017.
3. Malfertheiner P, Megraud F, O'Morain CA, Atherton J, Axon AT, Bazzoli F. European Helicobacter Study Group. Management of Helicobacter pylori infection - the Maastricht IV/Florence consensus report. *Gut*. 2012; 61: 646-664.
4. Chen TS, Chang FY. Clinical characteristics of Helicobacter pylori-negative duodenal ulcer disease. *Hepato-gastroenterology*. 2008; 55: 1615-1618.
5. Mohamed NA, Abd El Moez Azzam NF. Helicobacter pylori Associated Psychiatric Disorders among Bahraini Adult Presenting with .Function Dyspepsia. *J Med Sciences*. 2020; 20: 55-59.
6. Goodwin RD, Talley NJ, Hotopf M, Cowles RA, Galea S, Jacobi FA link between physician-diagnosed ulcer and anxiety disorders among adults. *Annals of epidemiology*. 2013; 23:189-192.
7. Agah S, Khedmat H, Ghamar-Chehred ME, Hadi R, Aghaei A. Female gender and Helicobacter pylori infection, the most important predisposition factors in a cohort of gastric cancer: A longitudinal study. *Caspian j Internal Med*. 7: 136; 2016.
8. Heijtz RD, Wang S, Anuar F, Qian Y, Björkholm B, Samuelsson A, Pettersson S. Normal gut microbiota modulates brain development and behavior. *Proceedings National Academy Sci*. 2011; 108: 3047-3052.
9. Al-Makdad AM, Al-Dholae MH, Thabet AAK, Al-Haimi MA, Balfaqih OS, Al-Hadad AM. Prevalence of Helicobacter pylori infection in Yemeni patients. *Yemeni Jou Med Sci*. 2013; 7: 33-38.
10. Almashhadany DA, Mayass SM. Prevalence of Helicobacter pylori in Human in Dhamar governorate/Yemen. *J Med Pharmaceutical Sci*. 2018; 2: 18-1.
11. Alyahawi A, Alkaf A, Alzaghroori S. Prevalence of helicobacter pylori among asymptomatic populations in Sana'a, Yemen. *Uni J Pharma Res*. 2018; 3: 31-35.
12. Budzyński J, Kłopotcka M. Brain-gut axis in the pathogenesis of Helicobacter pylori infection. *World Jou Gastroenterology: WJG*. 2014; 20: 5212.
13. Piriyapong K, Tangaroonsanti A, Mahachai V, Vilaichone RK. Helicobacter pylori infection impacts on functional dyspepsia in Thailand. *Asian Pac J Cancer Prev*. 2014; 15: 10887-91.
14. Mohamed DM, Elrassas H. Helicobacter Pylori Associated Depression among Patients Presenting with Epigastric Pain. *Egyptian J Hospital Med*. 2023; 90: 2315-2320.
15. Eshraghian A. Epidemiology of Helicobacter pylori infection among the healthy population in Iran and countries of the Eastern Mediterranean Region: a systematic review of prevalence and risk factors. *World j gastroenterol: WJG*. 2014; 20:17618.
16. Akeel M, Elmakki E, Shehata A, Elhafey A, Aboshouk T, Ageely H, et al. Prevalence and factors associated with *H. pylori* infection in Saudi patients with dyspepsia. *Electronic physic*. 2018; 10: 7279.



17. Marie MAM. Alimentary tract: Seroprevalence of helicobacter pylori infection in large series of patients in an urban area of Saudi Arabia. *J Korean Soci Gastroenterol*. 2008; 52: 226-229.
18. Gunaid AA, Hassan NA, Murray-Lyon I. Prevalence and risk factors for Helicobacter pylori infection among Yemeni dyspeptic patients. *Saudi Med J*. 2003; 24: 512-517.
19. Levenstein S, Kaplan GA, Smith MW. Psychological predictors of peptic ulcer incidence in the Alameda County Study. *J Clini Gastroenterol*. 1997; 24: 140-146.
20. Nguyen T, Ramsey D, Graham D, Shaib Y, Shiota S, Velez M, et al. The Prevalence of H elicobacter pylori Remains High in African American and Hispanic Veterans. *Helicobacter*. 2015; 20: 305-315.
21. Levenstein S. The very model of a modern etiology: a biopsychosocial view of peptic ulcer. *Psychosoma Med*. 2000; 62: 176-185.
22. Cader J, Domagala Z, Paradowski L, Rymaszewska J, Blonski W, et al. Is there any relation of Helicobacter pylori infection to anxiety and depressive symptoms? *Gastroenterologia Polska*. 2007; 14: 397.
23. Kabeer KK, Ananthakrishnan N, Anand C, Balasundaram S. Prevalence of Helicobacter pylori infection and stress, anxiety or depression in functional dyspepsia and outcome after appropriate intervention. *J Clinical Diagnos Resear: JCDR*. 2018; 11: VC11.
24. Sodhi JS, Javid G, Zargar SA, Tufail S, Shah A, Khan BA, et al. Prevalence of H elicobacter pylori infection and the effect of its eradication on symptoms of functional dyspepsia in Kashmir, India. *J Gastroenterol Hepatol*. 2013; 28: 808-813.
25. Tennant C, Goulston K, Langeluddecke P. Psychological correlates of gastric and duodenal ulcer disease. *Psychologi Med*. 1986; 16: 365-371.
26. Kivrak Y, Kokacya H, Copoglu US, Dokuyucu R, Ari M, Sulu B, et al. Does Helicobacter Pylori Cause Psychiatric Symptoms in Dyspeptic Patients? *Acta Medica*. 2014; 30: 1271.